arXiv:1409.6850v1 [cond-mat.stat-mech] 24 Sep 2014

Equilibrium statistics of an infinitely long chain in the severe confined geometry: Rigorous results

Pramod Kumar Mishra
Department of Physics, DSB Campus, Kumaun University
Nainital-263 002, India

Abstract : We analyze the equilibrium statistics of a long linear homopolymer chain confined in between two flat geometrical constraints under good solvent condition. The chain is occupying two dimensional space and geometrical constraints are two impenetrable lines for the two dimensional space. A fully directed self avoiding walk lattice model is used to derive analytical expression of the partition function for the given value of separation in between the impenetrable lines. The exact values of the critical exponents ($\nu_{||}, \nu_{\perp}, \nu$ and γ_1) were obtained for different value of separations in between the impenetrable lines. An exact expression of the grand canonical partition function of the confined semiflexible chain is also calculated for the given value of the constraints separation using generating function technique.

PACS Nos.: 05.70.Fh, 64.60 Ak, 05.50.+q, 68.18.Jk, 36.20.-r

1 Introduction:

The subject of confinement of a long linear polymer chain has received much attention in the past a few years due to advances in single molecule based experiments [1, 2, 3, 4, 5, 6], and also see references quoted therein. It is now possible to measure conformational properties of a confined polymer chain [7]. Such study can reveal wealth of useful information regarding elastic properties and molecular organization of the chain in the crowded geometry. It may also be useful in understanding molecular processes occurring in the living cells.

Lattice model of self-avoiding walk is extensively used to study conformational properties of a long linear polymer chain. A rich phase diagram of the surface interacting polymer chain has been reported by various authors [1, 2, 3, 4, 5, 6, 7, 8]. The values of critical exponents were also reported and these exponents were found useful in understanding thermodynamics of the chain. However, a few aspects of a confined long chain in the severe confined geometry still require attention. These studies may find useful ap-

plications in the steric stablisation of the colloids dispersions, surface coating and adsorption behaviour of the gels [1, 2, 3, 4].

In this paper we consider a long chain confined in between two impenetrable lines. The separation in between confining lines is varied in the unit of monomer size. An approach of canonical ensemble theory is used to derive exact expression of the canonical partition function of the confined flexible chain for a given value of separation (M) in-between the confining lines. The value of critical exponents $\nu_{||}, \nu_{\perp}, \nu$ and γ_1 were determined for the separation (M) in between the confining lines. We have also derived the exact expression of the grand canonical partition function of the confined semi-flexible chain using generating function technique.

The paper is organized as follows: In Sec. 2, lattice model of fully directed self avoiding walk is described for a confined linear polymer chain under good solvent condition on a square lattice. In sub-section 2.1, we use canonical ensemble theory to derive expression of the thermodynamic parameters of the confined single chain. Analytical expressions of the critical exponents were obtained for the confined flexible chain. In the subsection 2.2 the grand canonical ensemble theory is used to derive exact partition function of an infinitely long semiflexible chain for a given value of separation in between the confining impenetrable lines. Finally, in Sec. 3 we summarize and discuss the results obtained.

2 Model and method

A model of fully directed self-avoiding walks [9] on a square lattice is used to investigate the possibility of an adsorption transition of an infinitely long linear semiflexible homopolymer chain on geometrical constraints, when the chain is confined in between two impenetrable stair shaped surfaces under good solvent condition (as shown schematically in figure 1). The directed walk model is restrictive in the sense that the angle of bending has a unique value, that is 90° for a square lattice and directedness of the walk amounts to certain degree of stiffness in the walks of the chain because the different directions of the space are not treated equally. Since, the directed self avoiding walk model can be solved analytically and therefore gives exact values of the partition function of the polymer chain. We consider a fully directed self avoiding walk ($FDSAW$) model, therefore, the walker is allowed to take steps along $+x$, and $+y$ directions on a square lattice in between the constraints.

A fully directed self avoiding walk [1] is used to model a long linear



Figure 1: This schematic diagram shows a linear polymer chain confined in between two impenetrable lines. The lines are $y = 0$ and $y = 5$ *i. e.* the separation (M) in between the confining lines is 5 units. One end of the confined chain is grafted at (O) lower impenetrable line.

homo-polymer chain in between two impenetrable lines ($y = 0$ and $y = M$) on a square lattice under good solvent condition (as shown in figure-1, where $M = 5$). In other words, the separation is five units in-between impenetrable lines. One end of the confined chain is grafted on an impenetrable line $y = 0$ and the walker is allowed to take steps along $+x$ and $+y$ directions in between the two impenetrable lines. The spacing in-between two impenetrable lines is increased by one unit of monomer (step) size. Since, the walk is fully directed; therefore, it allows certain degree of restriction on the chain. However, directed walk model is analytically solvable [1, 3, 6, 8, 9, 10] and exact values of the critical exponents can be determined using this model.

The stiffness of the chain is accounted by introducing an energy $\epsilon_b(> 0)$ for each bend occurring in the chain. The stiffness weight is $k = exp(-\beta\epsilon_b)$, where $\beta = (k_B T)^{-1}$ is inverse of the temperature. If, $k = 1$ or $\epsilon_b = 0$, the chain is flexible and for $0 < k < 1$ or $0 < \epsilon_b < \infty$ the chain is semiflexible. However, if $\epsilon_b \to \infty$ or $k \to 0$, the polymer chain is like a rigid rod.

The generating function technique is used to derive exact expression of the partition function of a confined semiflexible chain, when the separation in between two impenetrable lines is M units.

2.1 A long linear chain confined in between the impenetrable lines: Canonical Ensemble Approach

We consider a long linear chain of N monomers confined in between two impenetrable lines. The lines are separated to each other by a distance of M units. The number of conformations $[C_N^M(N_x, N_y)]$ of the confined chain varies with number of steps N_x and N_y (where, N_x is the number of steps

along $+x$ direction and N_y is the number of steps along $+y$ direction) of a walk of N steps in between the confining lines. A list of values of the conformations are shown in the table -1 of a confined flexible chain of N monomers and the separation in between the confining impenetrable lines is M.

The total number of conformations $[Z_N^M]$ of the confined chain of N monomers variaes with separation M in between the impenetrable lines and its value for a flexible chain is written as,

$$Z_N^M = \sum_{N_y=0}^{M} C_N^M(N_x, N_y) = \sum_{K=1}^{M} \frac{\Pi_{P=1}^{K}[N-(P-1)]}{K!} + 1 \tag{1}$$

However, the number of suppressed conformations of the confined flexible chain of N monomers is $2^N - Z_N^M$, when the separation is M units in between the confining impenetrable lines.

An average value of the parallel component $(R_x(M))$ of the end-to-end distance $(R_E(M))$ is calculated using following relation,

$$< R_x^2(M) >= \frac{\sum_{N_y=0}^{M}(N-N_y)^2 C_N^M(N_x, N_y)}{Z_N^M} \tag{2}$$

where, $R_x(M) = N_x * a = (N - N_y) * a$, a is lattice parameter of a square lattice. The value of a is taken unity for the sake of mathematical simplicity.

In the long length limit of the confined flexible chain $R_x(M)$ scales as,

$$< R_x(M) >_{N\to\infty} \sim N(1 + \frac{M}{N}) \tag{3}$$

While, the average value of the perpendicular component $R_y(M)$ of the end-to-end distance of the confined chain is determined by following equation,

$$< R_y^2(M) >= \frac{\sum_{N_y=0}^{M} N_y^2 C_N^M(N_x, N_y)}{Z_N^M} \tag{4}$$

where, $R_y(M) = N_y * a$.

In the long length limit of the confined flexible chain $R_y(M)$ scales as,

$$< R_y(M) >_{N\to\infty} \sim M(1 + \frac{M}{N}) \tag{5}$$

The average value of end-to-end distance is,

Table 1: The number of conformations ($C_N^M(N_x, N_y)$) of a N step long confined flexible chain is shown in this table. The separation is M units in between the confining impenetrable lines.

N_x	$N_y(= N - N_x)$	$C_N^M(N_x, N_y)$
N-M	M	$\frac{N(N-1)(N-2)...[N-(M-1)]]}{M!}$
N-(M-1)	M-1	$\frac{N(N-1)(N-2)...[N-(M-2)]]}{(M-1)!}$
N-(M-2)	M-2	$\frac{N(N-1)(N-2)...[N-(M-3)]]}{(M-2)!}$
N-(M-3)	M-3	$\frac{N(N-1)(N-2)...[N-(M-4)]]}{(M-3)!}$
.	.	.
.	.	.
.	.	.
.	.	.
.	.	.
N-4	4	$\frac{N(N-1)(N-2)(N-3)}{4!}$
N-3	3	$\frac{N(N-1)(N-2)}{3!}$
N-2	2	$\frac{N(N-1)}{2!}$
N-1	1	N
N	0	1

$$< R_E^2(M) >= \frac{\sum_{N_y=0}^{M}[N_y^2 + (N - N_y)^2]C_N^M(N_x, N_y)}{Z_N^M} \tag{6}$$

The universal configurational exponent γ_1 of the confined flexible chain with one end grafted on an impenetrable line ($y = 0$) is calculated using formula,

$$\gamma_1(M) = \frac{Ln(\frac{Z_{N+P}^M Z_{N-P}^M}{(Z_N^M)^2})}{Ln(\frac{N^2-P^2}{N^2})} \tag{7}$$

where, N and P is the number of monomers in the confined flexible chain. We have found in the long length limit of the confined chain,

$$\frac{Z_{N+P}^M Z_{N-P}^M}{(Z_N^M)^2} = \frac{N^2 - P^2}{N^2} \tag{8}$$

The value of $\gamma_1(M)$=1.



Figure 2: It has been shown in this figure that the impenetrable lines (geometrical constraints) supress huge number of conformations of the confined flexible chain. When $N = 2M$, fifty percent conformations of the confined flexible chain is not polymerized due to presence of impenetrable lines.

2.2 *An infinitely long semiflexible chain in between two impenetrable lines: Grand Canonical Ensemble approach*

The grand canonical partition function of an infinitely long linear semiflexible homo-polymer chain confined in between two impenetrable lines (as shown in figure-1) is written as,

$$G(k) = \sum_{N=0}^{N=\infty} \sum_{all\ walks\ of\ N\ steps} g^N k^{N_b} \tag{9}$$

where, N_b is the total number of bends in a self avoiding walk of N steps (bonds or monomers), and g is step fugacity of the walker making steps in the space lying in-between the impenetrable lines.

The method of generating function technique [1] is used to calculate the grand canonical partition function of the confined semiflexible homopolymer chain (as shown in figure-1).

The components $[X_M(k,g)\&Y_M(k,g)]$ of the grand canonical partition function $G_M(k,g)$ of the confined semiflexible chain, when the separation in-between the confining impenetrable lines is equal to M units, can be written as,

$$X_M(k,g) = \frac{g}{1-g} + \frac{gk(1-g+gk)}{(1-g)^2}[u_1+u_2+u_3+u_4+u_5] \qquad (10)$$

where, $u_1 = \sum_{P=1}^{M} g^P + \frac{g^2k^2}{1-g}\sum_{P=1}^{M-1} Pg^P$

$u_2 = [\frac{g^2k^2}{1-g}]^2 \sum_{P=1}^{M-2} \frac{P(P+1)}{2!} g^P + [\frac{g^2k^2}{1-g}]^3 \sum_{P=1}^{M-3} \frac{P(P+1)(P+2)}{3!} g^P + \ldots$

$u_3 = [\frac{g^2k^2}{1-g}]^{M-3} \sum_{P=1}^{3} \frac{P(P+1)(P+2)\ldots(P+M-4)}{(M-3)!} g^P +$

$u_4 = [\frac{g^2k^2}{1-g}]^{M-2} \sum_{M=1}^{2} \frac{P(P+1)(P+2)+\ldots(P+M-3)}{(M-2)!} g^P +$

$u_5 = [\frac{g^2k^2}{1-g}]^{M-1} \sum_{P=1}^{1} \frac{P(P+1)(P+2)\ldots(P+M-2)}{(M-1)!} g^P$

and $P \leq M$ (and M=1,2,3,4...,∞) and $X_M(k,g)$ is the sum of the Boltzmann weight of all the walks having first step along $+x$-direction.

$$X_{M\to\infty}(k,g) = \frac{g}{1-g-gk} \qquad (11)$$

provided $g < \frac{1}{1+k}$ *i. e.* g_c [8].

Similarly, $Y_M(k,g)$ is the sum of Boltzmann weight of all the walks of the confined polymer chain having first step along $+y$ direction. It can be written as,

$$Y_M(k,g) = \frac{1-g+gk}{1-g}[v_1+v_2+v_3+v_4+v_5] \qquad (12)$$

where,

$v_1 = \sum_{P=1}^{M} g^P + \frac{g^2k^2}{1-g}\sum_{P=1}^{M-1} Pg^P +$

$v_2 = [\frac{g^2k^2}{1-g}]^2 \sum_{P=1}^{M-2} \frac{P(P+1)}{2!} g^P + [\frac{g^2k^2}{1-g}]^3 \sum_{P=1}^{M-3} \frac{P(P+1)(P+2)}{3!} g^P + \ldots$

$v_3 = [\frac{g^2k^2}{1-g}]^{M-3} \sum_{P=1}^{3} \frac{P(P+1)(P+2)\ldots(P+M-4)}{(M-3)!} g^P +$

$v_4 = [\frac{g^2k^2}{1-g}]^{M-2} \sum_{M=1}^{2} \frac{P(P+1)(P+2)+\ldots(P+M-3)}{(M-2)!} g^P +$

$v_5 = [\frac{g^2k^2}{1-g}]^{M-1} \sum_{P=1}^{1} \frac{P(P+1)(P+2)\ldots(P+M-2)}{(M-1)!} g^P]$

and,

$$Y_{M\to\infty}(k,g) = \frac{g}{1-g-gk} \tag{13}$$

for $g < g_c$. The grand canonical partition function ($Z_M(k,g)$) of the confined chain can be calculated by summing the above two components of the partition function,

$$Z_M(k,g) = \frac{g}{1-g} + [\frac{1-g+gk}{1-g}]^2[w_1+w_2+w_3+w_4+w_5] \tag{14}$$

where,
$w_1 = \sum_{P=1}^{M} g^P + \frac{g^2k^2}{1-g}\sum_{P=1}^{M-1} Pg^P +$
$w_2 = [\frac{g^2k^2}{1-g}]^2 \sum_{P=1}^{M-2} \frac{P(P+1)}{2!}g^P + [\frac{g^2k^2}{1-g}]^3 \sum_{P=1}^{M-3} \frac{P(P+1)(P+2)}{3!}g^P + \ldots$
$w_3 = [\frac{g^2k^2}{1-g}]^{M-3} \sum_{P=1}^{3} \frac{P(P+1)(P+2)...(P+M-4)}{(M-3)!}g^P +$
$w_4 = [\frac{g^2k^2}{1-g}]^{M-2} \sum_{M=1}^{2} \frac{P(P+1)(P+2)+...(P+M-3)}{(M-2)!}g^P +$
$w_5 = [\frac{g^2k^2}{1-g}]^{M-1} \sum_{P=1}^{1} \frac{P(P+1)(P+2)...(P+M-2)}{(M-1)!}g^P]$
and substituting $M \to \infty$ in above equation of the partition function, we recover $Z_\infty(k,g) = \frac{2g}{1-g-gk}$ [8], provided $g < g_c(= \frac{1}{1+k})$ [8].

3 Result and discussion

A fully directed self avoiding walk lattice model is used to calculate canonical partition function of a long linear flexible polymer chain confined in between two impenetrable lines (as shown in figure-1). The confined chain is uccupying space in between two confining lines and the chain is under good solvent condition. The separation (M) in-between the confining impenetrable lines is increased in the unit of monomer (step) size. The exact expression of the canonical partition function of the confined flexible chain is obtained for a given value of the separation in between the confining lines. The critical value of exponents were determined and the exact expression of the grand canonical partition function of a confined semiflexible chain is also derived. We have chosen monomer (step) size unity for the sake of mathematical simplicity.

We have shown the variation in the number ($C_N^M(N_x,N_y)$) of conformations (entropy) of the confined flexible chain of N monomers for a given separation (M) in between the confining lines (as shown in table-1). The effect of confinement on the statistics of a flexible chain is illustrated by the

figure 2. It has been found that the 50% conformations of the confined flexible chain are suppressed due to confinement, when $N = 2M$. The number of suppressed conformations is $2^N - Z_N^M$ of a confined flexible chain of N monomers confined in between two impenetrable lines separated by M units.

The average value of end-to-end distance of the confined flexible polymer chain is modified due to confinement. The average value of the parallel component ($R_x(M)$) of the end-to-end distance $[R_E(M)]$ and $R_E(M)$ of the confined flexible chain scales as, $< R_x > \sim N(1 + \frac{M}{N})$ and the perpendicular component of end-to-end distance ($R_y(M)$) scales with the confined chain size as $< R_y > \sim M(1 + \frac{M}{N})$. Therefore, the value of $\nu_{||} = 1$ and $\nu = 1$ while, $\nu_\perp = 0$ for the confined flexible chain, provided the value of M is finite. It has been concluded from the value of critical exponent ν and $\nu_{||}$ that the confined chain in quasi-one dimension. Therefore, value of $\gamma_1(M)$ is unity for a flexible chain confined in-between two parallel impenetrable lines.

It has also been found that the critical value of step fugacity (g_c) of the confined flexible chain is 1, when the separation in-between the confining impenetrable lines is M. However, $g_c = \frac{1}{1+k}$[8] as the value of $M \to \infty$.

3.1 Acknowledgements

Author thankfully acknowledge financial support received from Department of Science and Technology, New Delhi (Project no. SR/FTP/PS-122/2010).

References

[1] V Privman, G Forgacs and H L Frisch 1988 Phys. Rev. B **37** 9897; Privman V and Svrakic N M 1989 *Directed Models of Polymers, Interfaces, and Clusters: Scaling and Finite-Size Properties* (Springer, Berlin).

[2] J des Cloiseaux and G Jannink 1990 *Polymers in Solution* (Clarendon, Oxford); E Eisenriegler 1993 *Polymers Near Surfaces* (World Scientific, Singapore).

[3] C Vanderzande 1998 *Lattice Models of Polymers* (Cambridge University Press, UK).

[4] *Structure and dynamics of confined polymers* 2002 (ed. J J Kasianowicz *et. al*, Kluwer Academic Publishesr, Dordrecht).

[5] P K Mishra and S. Kumar 2004 J. Chem. Phys. **121** 8642; P Cifra and T Bleha 2010 Euro. Phys. J. E **32** 273; P Cifra 2009 J. Chem. Phys. **131** 224903.

[6] R Brak, G K Iliev, A L Owczarek and S G Whittington J. Phys A: Math. Theor. 2010 **43** 135001; R Brak, A L Owcarek, A Rechnitzer and S G Whittington 2005 J. Phys. A **38** 4309; Erich D Bain, Salomom T-Cohen and J Genzer 2013 Macromol. Theory Simul. **22** 8.

[7] W D Volkmuth and R H Austin 1992 Nature **358** 600; W D Volkmuth, T Duke, M C Wu, R H Austin and A Szabo 1994 Phys. Rev. Lett. **72** 2117.

[8] P K Mishra, S Kumar and Y Singh 2003 Physica A **323** 453.

[9] P K Mishra, 2010 J. Phys.: Cond. Matt. **22** 155103; P K Mishra, 2010 Phase Tran. **83** 47.

[10] P K Mishra, 2011 Phase Tran. **84** 291; P K Mishra 2014 Cond. Matt. Phys. **17(2)** 23001.